

## The Flora of Upper Ceyhan Valley (Kahramanmaraş/Turkey)

Adem TATLI

Dumlupınar University, Art & Science Faculty, Biology Dept. Kütahya - TURKEY

Hasan AKAN

Harran University, Art & Science Faculty, Biology Dept. Şanlıurfa- TURKEY

A. Zafer TEL

Dumlupınar University, Art & Science Faculty, Biology Dept. Kütahya - TURKEY

Cemil KARA

Sütçü İmam University, Art & Science Faculty, Biology Dept. K.Maraş- TURKEY

Received: 22.11.2000

Accepted: 11.01.2002

**Abstract:** This investigation was carried out between 1994 and 1995 in order to determine the flora of Upper Ceyhan Valley. The research area is situated in the province of Kahramanmaraş and within C6 of the grid system. In the area, 412 taxa belonging to 68 families and 262 genera were determined. Twenty-eight (6.7%) taxa are endemics for Turkey. In the distribution of the taxa according to the floristic regions, Mediterranean elements with a rate of 23.5% are the most common. Irano-Turanian elements with a rate of 17.2% and Euro-Siberian elements with a rate of 3.4% were found. The families which have the most taxa in the research area are *Fabaceae* (60 spp.), *Asteraceae* (52 spp.), *Poaceae* (31 spp.), *Brassicaceae* (24 spp.) and *Lamiaceae* (20 spp.). Concerning the number of taxa, the major genera in this region are *Trifolium* L. (10 spp.), *Vicia* L. (9 spp.), *Euphorbia* L. (8 spp.), *Papaver* L. (6 spp.), *Galium* L. (6 spp.) and *Astragalus* L. (4 spp.).

**Key Words:** Flora, Upper Ceyhan Valley, Kahramanmaraş.

### Yukarı Ceyhan Vadisi Florası (Kahramanmaraş/Türkiye)

**Özet:** Bu araştırma, 1994-1995 yılları arasında Yukarı Ceyhan Vadisi florasını tespit etmek üzere yapılmıştır. Araştırma alanı, Kahramanmaraş il sınırları içerisinde ve Grid sistemine göre C6 karesindedir. Araştırma bölgemizde, 68 familya ve 262 cinsle ait 412 takson tespit edildi. Bunlardan, 28 (%6,7) takson Türkiye için endemiktir. Bitkilerin fitocoğrafik bölgelere dağılımında; %23,5 oranı ile Akdeniz elementleri ilk sırayı almaktadır. İran-Turan elementleri %17,2 ve Avrupa-Sibirya elementleri ise %3,4 oranlarıyla temsil edilir. Sahada en çok taksona sahip olan familyalar; *Fabaceae* (60 tür), *Asteraceae* (52 tür), *Poaceae* (31 tür), *Brassicaceae* (24 tür) ve *Lamiaceae* (20 tür)'dır. Alanda en çok taksona sahip olan cinslerin sıralanışı ise şöyledir: *Trifolium* L. (10 tür), *Vicia* L. (9 tür), *Euphorbia* L. (8 tür), *Papaver* L. (6 tür), *Galium* L. (6 tür) ve *Astragalus* L. (4 tür)'dur.

**Anahtar Sözcükler:** Flora, Yukarı Ceyhan Nehir Vadisi, Kahramanmaraş.

### Introduction

This research was carried out to determine the flora of Upper Ceyhan Valley in Kahramanmaraş. The vegetation of the area has been extensively destroyed by anthropogenic effects and, therefore, plants can barely survive under the unsuitable agricultural conditions.

Close to our research area, some floristic and vegetation studies were carried out by Akman (1973), Çetik (1981), Duman (1991, 1995), Düzenli (1976), Eyce (1987), Eyce and Ocakverdi (1987), Quezel (1973), Quezel and Pamukçuoğlu (1973), Uslu (1977), Vural et al. (1999) and Yıldız (1984).

Balls, Davis, Guichard, Haradjian, Haussknecht, Post and Sintenis have made some collections around

Kahramanmaraş. Among them, especially Balls, Davis and Haussknecht have some collections from the Ceyhan valley (Davis 1965-1985).

We chose the research area, Upper Ceyhan Valley, for the following reasons:

- 1- It is between the Mediterranean and Irano-Turanian phytogeographic areas.
- 2- It was not studied previously in terms of flora and vegetation.
- 3- It is on the Anatolian Diagonal.
- 4- According to Zohary (1973), endemic plants are concentrated in this region.

In light of this information, our main goals may be summarized as follows:

- 1- to determine the flora of Upper Ceyhan Valley.
- 2- to determine rare and endemic plants of the research area. The research area is between Kılavuzlu Village and Menzelet Dam, which will be permanently under water.
- 3- to determine the main peculiarities of the vegetation types in the region.

#### Geographical Location


The research area is located between Mediterranean and Irano-Turanian phytogeographic regions and at the south of the Anatolian Diagonal (Zohary, 1973). The research area is situated in the C6 square and completely located within the province of Kahramanmaraş. The altitude of the area is between 500 and 1250 m.

Ceyhan River is fed by Aksu River, Bertiz Creek, Cemrengec Stream and Tekir Stream. The river originates at Elbistan and flows into the Mediterranean Sea.

There are 13 villages and a town in and around the research area. Generally, settlements are dense close to the river. The villages in and around the research area are Bulutoğlu, Çakırdere, Çavgalar, Demirlik, Ekizler, Elsen, Hacınınoğlu, Kamişlı, Kertel, Kılavuzlu, Koçlar, Payam and Sarıçukur. Baydemirli Village became a new town recently.

To the north of the research area are Berit Mountain, Bezirgan Mountain, Karadiken Mountain and Kişi Mountain; to the east, Engizek Mountain; to the south, Ahır Mountain, and to the west, Ali Hill (Figure 1).

Figure 1. The geographic map of study area



**Climate:** The natural flora and vegetation types and the meteorological data indicate that the research area is under the influence of the Mediterranean climate, which has rainfall in cold and relatively cold seasons, a dry season in summer and a maximum temperature of more than 25°C (Akman, 1990). Macchie plants such as *Pinus brutia* Ten., *Quercus coccifera* L., and *Arbutus andrachne* L., which show a natural spread in the research area, are indicators of a prevailing Mediterranean climate.

The annual average of temperature is 16.5°C and the total rain amount is 662.2 mm in Kahramanmaraş. The aridity period is between April and May. The precipitation regime is winter-spring-autumn-summer in Kahramanmaraş and Göksun (Meteoroloji Bülteni, 1994).

## Materials and Methods

To constitute research material, c. 700 plant specimens were gathered from Upper Ceyhan Valley by Cemil Kara between 1994 and 1995 during different vegetation seasons. A collector number was given and the specimens were dried according to herbarium methods. The plant samples collected from the region were kept in the herbarium of Dumlupınar University, Kütahya, and of Kahramanmaraş Sütçü İmam University. "Flora of Turkey and the East Aegean Islands" [(Davis, 1965-1985, Davis et al., 1988)] was used to identify plant samples.

The list of plants is set out according to the order in the *Flora of Turkey*. In the floristic list of this article, the following details are given: family name, species name and author(s), the name of district, habitat of plant, altitude, collection date, collector number given by C. Kara, endemism and the phytogeographic element.

After checking the *Flora of Turkey*, the taxa that we were unable to collect during this study but had been collected by Davis, Balls and Haussknecht were added to the floristic list.

### The following abbreviations are used in the text:

Brj.:	Barajı
Boğ.:	Boğazi
E:	East
Euro-Sib.:	Euro-Siberian
HES:	Hydro Electric Power Station

Ir.-Tur :	Irano-Turanian
K.:	C. Kara
Ky:	Köyü
m:	meter
Medit:	Mediterranean
S:	South
subsp.:	subspecies
var.:	variety
W:	West

## Vegetation

The vegetational characteristics of the research area are generally as follows: at 500-600 m between Kılavuzlu village and the Menzelet dam, there are *Tamarix parviflora* DC., *Platanus orientalis* L., *Juniperus oxycedrus* L. subsp. *oxycedrus*, *Quercus cerris* L. var. *cerris* and *Quercus coccifera*. There are *Pinus brutia* forests around the Menzelet dam.

Under the forest, *Juniperus oxycedrus* subsp. *oxycedrus* and *Arbutus andrachne* which are Mediterranean elements, are spread at intervals.

In the fields, along the river *Adonis flammea* Jacq., *Anemone coronaria* L., *Anemone blanda* Schott & Kotschy and *Papaver rhoeas* L. can occasionally be found.

At the shallow parts of the river, there are *Juncus inflexus* L., *Juncus articulatus* L., *Plantago major* L., *Plantago lanceolata* L., *Rumex pulcher* L., *Sambucus nigra* L., *Equisetum hyemale* L., *E. arvensis* L. and *Adiantum capillus-veneris* L.

Along the riverside, *Salix pentandra* L. and *Populus nigra* L. subsp. *nigra* are vanguard trees. Around the erosion area of Kılavuzlu village, near the river, *Rhus coriaria* L. is a dominant group. In addition, with *Styrax officinalis* L. also *Capparis spinosa* L., which are adapted to the river's rocky parts, can be observed.

Between Çakırdere village and Kısık channel, along the riverside, *Rheum ribes* L. is spread at intervals.

In the *Pinus brutia* forests, near the river, *Onobrychis tournefortii* (Willd.) Desv., *Moluccella leavis* L., *Astragalus vaginans* DC., *A. gummifer* Labill. and *A. hamosus* L. are spread between Kılavuzlu village and the Güvercinlik dam.

In fields and roadsides, *Paliurus spina-christi* Mill., *Cerasus avium* (L.) Moench, *C. microcarpa* (C.A. Meyer) Boiss., *Reseda lutea* L., *Tribulus terrestris* L., *Centeurea iberica* Trevir. ex Sprengel, *Buglossoides arvensis* (L.) Johnston, *Verbena officinalis* L., *Poa bulbosa* L., *Veronica anagallis-aquatica* L., *Crupina crupinastrum* (Moris) Vis., *Stachys cretica* L., *Plantago major* L., *P. lanseolata* L.,

*Ononis natrix* L., *Trifolium campestre* Schreb., *Cardaria draba* (L.) Desv., *Medicago x varia* Martyn and *Vicia sericocarpa* Fenzl are observed.

In the regions appropriate for agriculture, some fruits and vegetables are grown, such as *Junglans regia* L., *Cerasus avium* (L.) Moench, *Vitis vinifera* L., *Morus nigra* L. and *Ficus carica* L.

#### THE FLORISTIC LIST

##### DIVISIO: PTERIDOPHYTA

###### 1- EQUISETACEAE

*Equisetum hyemale* L.

Kilavuzlu, roadsides, 500-600 m, 25.02.1995, K. 781.

*E. arvense* L.

Ceyhan HES, 500-600 m, 20.05.1995, K. 780.

*E. ramosissimum* Desf.

Kertel, streamside, 500-600 m, 08.08.1995, K. 607.

###### 2- ADIANTACEAE

*Adiantum capillus-veneris* L.

Ceyhan HES, 500-550 m, 08.04.1995, K. 782.

##### DIVISIO: SPERMATOPHYTA

###### SUBDIVISIO: GYMNOSPERMAE

###### 3- PINACEAE

*Pinus brutia* Ten.

Kertel to Menzelet, 600-800 m, 03.05.1995, K. 783.

Medit. element.

###### 4- CUPRESSACEAE

*Cupressus sempervirens* L.

Ceyhan HES, 500-600 m, 20.05.1995, K. 785.

*Juniperus oxycedrus* L. subsp. *oxycedrus*

Çakırdere, forest clearings, 800-900 m, 03.09.1995, K. 633.

###### 5- EPHEDRACEAE

*Ephedra campylopoda* C.A.Mey.

Güvercinlik, 500-600 m, 07.11.1995, K. 791.

Medit. element.

###### SUBDIVISIO: ANGIOSPERMAE

###### Classis: DICOTYLEDONES

###### 6- RANUNCULACEAE

*Delphinium peregrinum* L.

Kertel to Güvercinlik, 500-600 m, 27.04.1995, K. 326.

Medit. element.

*D. virgatum* Poir.

Kertel to Ceyhan HES, fields, 500-600 m, 20.08.1995, K. 609.

*Anemone blanda* Schott & Kotschy

Bulutoğlu, fields, c. 600 m, 11.03.1995, K. 324.

*A. coronaria* L.

Çalidere, fields, 800-900 m, 18.03.1995, K. 322.

Medit. element.

*Clematis orientalis* L.

Kertel, 500-600 m, 08.08.1995, K. 622.

*Adonis flammea* Jacq.

Kilavuzlu, 500-600 m, 20.04.1995, K. 320.

*A. annua* L.

Kilavuzlu to Bulutoğlu, fields, 500-600 m, 20.04.1995, K. 341.

*Ranunculus arvensis* L.

Kertel to Güvercinlik, fields, 500-600 m, 27.04.1995, K. 332.

*R. pinardii* (Stev.) Boiss.

Kertel to Güvercinlik, fields, 500-600 m, 27.04.1995, K. 327.

Ir.-Tur. element.

*Ceratocephalus falcatus* (L.) Pers.

Ceyhan HES, fields, c. 600 m, 08.04.1995, K. 323.

*Thalictrum orientale* Boiss.

Kertel, forest clearings, 500-600 m, 27.04.1995, K. 344.

Medit. element.

###### 7- PAPAVERACEAE

*Glaucium corniculatum* (L.) Rudolph subsp. *refractum* (Nábélek) Cullen

Güvercinlik, fields, 500-600 m, 20.05.1995, K. 440.

Endemic, Ir.-Tur. element.

*Glaucium acutidentatum* Hausskn. & Bornm.

Ceyhan HES, sandy fields, 500-600 m, 20.04.1995, K. 437.

Endemic, Ir.-Tur. element.

*Papaver glaucum* Boiss. & Hausskn.

Çakırdere to Kısık, 900-1000 m, 03.05.1995, K. 447.

Ir.-Tur element.

*P. rhoeas* L.

Ceyhan HES, fields, 500-600 m, 20.04.1995, K. 438.

*P. syriacum* Boiss. & Blanche

Güvercinlik, sandy places, 500-600 m, 20.05.1995, K. 442.

Medit. element.

*P. dubium* L.

S of Kertel, fields, 500-600 m, 20.05.1995, K. 450.

*P. clavatum* Boiss. & Hausskn.

S of Kertel, fields, 500-600 m, 20.04.1995, K. 439.

Medit. element.

*P. argemone* L.

Bulutoğlu, fields, 500-600 m, 20.04.1995, K. 439.

Medit. element.

<i>Fumaria asepala</i> Boiss.	18.03.1995, K. 242.	10-RESEDACEAE
Bulutoğlu, fields, 08.04.1995, K. 564.	Alyssum huetii Boiss.	<i>Reseda lutea</i> L. var. <i>lutea</i>
Ir.-Tur. element.	Ceyhan HES, roadside, 500-600 m, 11.03.1995, K. 235.	Sarıçukur to Kamişlı, fields, c. 700 m, 17.06.1995, K. 776.
8-BRASSICACEAE	Endemic, Ir.-Tur. element.	11-CISTACEAE
<i>Sinapis arvensis</i> L.	<i>Alyssum minus</i> (L.) Rothm. var. <i>minus</i>	<i>Helianthemum salicifolium</i> (L.) Mill.
S of Kertel, landslide places, 500-600 m, 08.04.1995, K. 253/a.	Koçlar to Kamişlar, fields, c. 1000 m, K. 231.	Kertel to Güvercinlik, riverside, 500-600 m, 11.03.1995, K. 734.
<i>Raphanus raphanistrum</i> L.	<i>Clypeola jonthlaspi</i> L.	12-VIOLACEAE
S of Kertel, landslide places, c. 600 m, 08.04.1995, K. 253/b.	S of Kertel, landslide places, 500-600 m, 08.05.1995, K. 251.	<i>Viola modesta</i> Fenzl
<i>Crambe orientalis</i> L. var. <i>orientalis</i>	<i>Erophila minima</i> C.A.Mey.	Baydemirli, hills, 800-1000 m, 18.03.1995, K. 650.
W of Kertel, c. 600 m, 27.04.1995, K. 224.	S of Kertel, under forest, 500-600 m, 25.02.1995, K. 205.	13-PORTULACACEAE
Ir.-Tur. element.	Medit. element.	<i>Portulaca oleracea</i> L.
<i>Lepidium perfoliatum</i> L.	<i>Arabis aucheri</i> Boiss.	Ceyhan HES, fields, 500-600 m, 20.05.1995, K. 779.
Çakırdere, field clearings, 900-1000 m, 18.03.1995, K. 240.	S of Kertel, forest clearings, 500-600 m, 22.03.1995, K. 218.	14-CARYOPHYLLACEAE
<i>Cardaria draba</i> (L.) Desv. subsp. <i>draba</i>	<i>Cardamine hirsuta</i> L.	<i>Minuartia hirsuta</i> (M.Bieb.) Hand.-Mazz. subsp. <i>falcata</i> (Gris.) Mattf.
Kılavuzlu, meadow, 500-600 m, 05.04.1995, K. 206.	S of Kertel, forest clearings, 500-600 m, 22.03.1995, K. 215.	Güvercinlik to Menzelet, fields, 500-600 m, 20.04.1995, K. 483.
<i>Iberis taurica</i> DC.	<i>Matthiola longipetala</i> (C.A.Venturi) DC. subsp. <i>longipetala</i>	<i>M. sclerantha</i> (C.E.C.Fisch. & C.A.Mey.) Thell.
Koçlar, fields, 900-1000 m, 03.04.1995, K. 219.	Baydemirli, hill slopes, 900-1000 m, 18.03.1995, K. 723.	Çakırdere to Aliyakası, fields, 800-1000 m, 03.05.1995, K. 760.
<i>Aethionema arabicum</i> (L.) Andrz.	Ir.-Tur. element.	Ir.-Tur. element.
Kısık, fields, 1000 m, 18.03.1995, K. 223.	<i>Erysimum goniocaulon</i> Boiss.	<i>Stellaria media</i> (L.) Vill. subsp. <i>postii</i> Holmboe
<i>Thlaspi perfoliatum</i> L.	Bulutoğlu to Ceyhan HES, roadside, 500-600 m, 12.06.1995, K. 256.	S of Kılavuzlu and Kertel, hills, 500-600 m, 27.04.1995, K. 472.
S of Kertel, under <i>Quercus coccifera</i> , 500- 600 m, 25.02.1995, K. 209.	Medit. element.	<i>Cerastium perfoliatum</i> L.
<i>Capsella rubella</i> Reut.	<i>Erysimum repandum</i> L.	Kılavuzlu, sandy places, 500-600 m, 27.04.1995, K. 475.
Güvercinlik, roadside, 500-600 m, 13.03.1995, K. 208.	Koçlar, fields, 800-1000 m, 18.03.1995, K. 225.	<i>C. brachypetalum</i> Pers. subsp. <i>roeseri</i> (Boiss. & Heldr.) Nyman
Medit. element.	<i>Camelina laxa</i> C.A.Mey.	Kılavuzlu, sandy places, 500-600 m, 27.04.1995, K. 481.
<i>Myagrum perfoliatum</i> L.	Kılavuzlu, landslide places, 500-600 m, 27.04.1995, K. 225.	<i>Holosteum umbellatum</i> L. var. <i>umbellatum</i>
Çakırdere, spaces of <i>Quercus</i> sp., 900 m, 3.05.1995, K. 213.	<i>Camelina rumelica</i> Velen	Kılavuzlu, forest clearings, 500-600 m, 11.03.1995, K. 471.
<i>Neslia apiculata</i> M.A.Fisch., C.A.Mey. & Avé- Lall.	Kılavuzlu, landslide places, 500-600 m, 27.04.1995, K. 212.	<i>Sagina apetala</i> Ardré
S of Kertel, landslide places, 500-600 m, 08.04.1995, K. 250.	9-CAPPARACEAE	Kılavuzlu, roadside, 500-600 m, 11.03.1995, K. 473.
<i>Peltaria angustifolia</i> DC.	<i>Capparis spinosa</i> L. var. <i>spinosa</i>	
S of Kertel, under forest, 500-600 m, 13.03.1995, K. 207.	Bulutoğlu, roadside, 500-600 m, 20.06.1995, K. 666.	
Medit. element.	Medit. element.	
<i>Alyssum linifolium</i> Steph. ex Willd.		
Baydemirli, fields, 900-1000 m,		

<i>Loeflingia hispanica</i> L.	<i>Polygonum salicifolium</i> Brouss. ex Willd.	<i>L. virgultorum</i> Boiss. & Heldr.
Kılavuzlu, riverside, 500-600 m, 08.04.1995, K. 487.	Kertel, rocky places, 500-600 m, 03.11.1995, K. 668.	S of Kertel, forest, 500-600 m, 05.06.1995, K. 545.
<i>Petrorhagia velutina</i> (Guss.) P.W.Ball & Heywood	<i>Rumex pulcher</i> L.	Medit. element.
Sarıçukur, fields, 700-800 m, 17.06.1995, K. 478.	Kertel to Güvercinlik, riverside, 500-600 m, 05.06.1995, K. 668.	<b>22-GERANIACEAE</b>
<i>Velezia rigida</i> L.	<b>17-CHENOPodiACEAE</b>	<i>Geranium lucidum</i> L.
Kamışlı, riverside, 700-800 m, 03.05.1995, K. 760.	<i>Chenopodium botrys</i> L.	S of Kertel, riverside, 500-600 m, 22.03.1995, K. 489.
<i>Saponaria glutinosa</i> M.Bieb.	Kertel to Güvercinlik, riverside, 500-600 m, 05.06.1995, K. 767.	<i>G. rotundifolium</i> L.
Kılavuzlu, sandy places, 500-600 m, 25.02.1995, K. 475.	<b>18-TAMARICACEAE</b>	Ceyhan HES to Güvercinlik, riverside, 500-600 m, 11.03.1995, K. 491.
<i>Gypsophila hispida</i> Boiss.	<i>Tamarix parviflora</i> DC.	<i>G. molle</i> L. subsp. <i>molle</i>
Ceyhan HES to Güvercinlik, roadside, 500-600 m, 08.08.1995, K. 594.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 05.06.1995, K. 674.	Hacıminoğlu to Demirlik, riverside, 900-1100 m, 11.03.1995, K. 500.
Ir.-Tur. element.	<b>19-HYPERICACEAE</b>	<i>G. columbinum</i> L.
<i>Vaccaria pyramidata</i> Medik. var. <i>grandiflora</i> (C.E.C.Fisch. ex DC.) Cullen	<i>Hypericum lydium</i> Boiss.	Bulutoğlu, riverside, 500-600 m, 08.04.1995, K. 497.
Ceyhan HES, fields, 500-600 m, 20.04.1995, K. 476.	Menzelet to Sarıcak, hills, 600-800 m, 17.06.1995, K. 751.	<i>G. stepporum</i> P.H.Davis
<i>Silene nuncupanda</i> Coode & Cullen	<b>20-MALVACEAE</b>	Kertel, riverside, 500-600 m, 22.03.1995, K. 490.
Sarıçukur to Kamışlı, hills, 800-1000 m, 17.06.1995, K. 477.	<i>Malva nicaeensis</i> All.	<i>Erodium botrys</i> (Cav.) Bertol.
Endemic.	Bulutoğlu, 500-600 m, 20.04.1995, K. 663.	Ceyhan HES, riverside, 500-600 m, 20.04.1995, K. 499.
<i>S. crassipes</i> Fenzl	<i>M. parviflora</i> L.	Medit. element.
Kılavuzlu, forest, 500-600 m, 22.03.1995, K. 470.	Bulutoğlu, roadside, 500-600 m, 20.05.1995, K. 660.	<i>E. malacoides</i> (L.) L'Hér.
<i>S. dichotoma</i> Ehrh. subsp. <i>dichotoma</i>	<i>Alcea striata</i> (DC.) Alef. subsp. <i>rufescens</i> (Boiss.) Cullen	Kılavuzlu, roadside, 500-600 m, 11.03.1995, K. 491.
Kısık, hills, 900-1000 m, 03.05.1995, K. 480.	Güvercinlik to Kertel, roadside, 500-600 m, 05.06.1995, K. 662.	Medit. element.
<b>15-ILLECEBRACEAE</b>	<i>A. apterocarpa</i> (Fenzl) Boiss.	<i>E. acaule</i> (L.) Bech. & Thell.
<i>Paronychia kurdica</i> Boiss. subsp. <i>kurdica</i> var. <i>kurdica</i>	Sarıçukur to Koçlar, fields, 600-900 m, 17.06.1995, K. 661.	Kılavuzlu, riverside, 500-600 m, 31.11.1994, K. 493.
Kılavuzlu, landslide places, 500-600 m, 22.03.1995, K. 726.	Endemic, Ir.-Tur. element.	Medit. element.
<b>16-POLYGONACEAE</b>	<b>21-LINACEAE</b>	<b>23-ZYGOPHYLLACEAE</b>
<i>Atraphaxis billardieri</i> Jaub. & Spach var. <i>billardieri</i>	<i>Linum flavum</i> L. subsp. <i>scabrinerve</i> (P.H. Davis) P.H. Davis	<i>Tribulus terrestris</i> L.
Kertel to Güvercinlik, hills, 500-600 m, 05.06.1995, K. 765.	S of Kertel, forest, 500-600 m, 05.06.1995, K. 781.	Ceyhan HES to Bulutoğlu, roadside, 500-600 m, 05.06.1995, K. 547.
Ir.-Tur. element.	Endemic, Ir.-Tur. element.	<b>24- VITACEAE</b>
<i>Rheum ribes</i> L.	<i>L. olympicum</i> Boiss.	<i>Vitis sylvestris</i> C.C.Gmel.
Çakırdere, riverside, 700-900 m, 03.05.1995, K. 533.	Ceyhan HES to Güvercinlik dam, 500-600 m, 20.05.1995, K. 778.	Kılavuzlu to Kertel, riverside, 500-600 m, 10.04.1995, K. 664.
Ir.-Tur. element.	Endemic.	<b>25-RHAMNACEAE</b>
		<i>Paliurus spina-christi</i> Mill.
		Kertel to Güvercinlik, 500-600 m, 20.05.1995, K. 714.

**26-ANACARDIACEAE**

*Rhus coriaria* L.  
Kılavuzlu to Güvercinlik, 500-600 m, 20.05.1995, K. 659.  
Medit. element.

*Pistacia vera* L.  
Bulutoğlu to Güvercinlik, 500-600 m, 05.06.1995, K. 657.  
Ir.-Tur. element.

*P. terebinthus* L. subsp. *palaestina* (Boiss.) Engl.  
Kılavuzlu to Kertel, 500-600 m, 20.06.1995, K. 658.  
Medit. element.

**27-FABACEAE**

*Astragalus hamosus* L.  
Kılavuzlu to Ceyhan HES, 500-600 m, 20.05.1995, K. 102.  
*A. gummifer* Labill.  
Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 06.07.1995, K. 71.  
Ir.-Tur. element.

*A. andrachneifolius* Fenzl  
Kamışlı to Baydemirli, riverside, 900-1000 m, 03.05.1995, K. 32.  
*A. vaginans* DC.  
Sarıçukur, forest clearings, 800-900 m, 17.06.1995, K. 5.  
Endemic.

*Glycyrrhiza aspera* Pal.  
Jehan (Ceyhan) valley, 730 m, Balls 1107.  
*Psoralea bituminosa* L.  
Bulutoğlu to Ceyhan HES, roadside, 500-600 m, 20.04.1995, K. 93.

Medit. element.

*Cicer pinnatifidum* Jaub. & Spach  
Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 20.05.1995, K. 3, 15 km N of Maraş, 500 m, D. 27464.

*Vicia villosa* Rothm. subsp. *villosa*  
Kertel to Güvercinlik, riverside, 500-600 m, 27.04.1995, K. 48.  
*V. cypria* Kotschy ex Unger & Kotschy  
Kılavuzlu, landslide places, 500-600 m, 25.02.1995, K. 115.

*V. hirsuta* (L.) Gray  
Kertel to Güvercinlik, riverside, 500-600 m, 27.04.1995, K. 46.  
*V. noeana* Reut. ex Boiss. var. *noeana*  
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.05.1995, K. 96.  
Ir.-Tur. element.

*V. peregrina* L.  
S of Kertel, riverside, 500-600 m, 22.03.1995, K. 83.

*V. sericocarpa* Fenzl var. *sericocarpa*  
S of Kertel, riverside, 500-600 m, 22.03.1995, K. 81.

*V. hybrida* L.  
S of Kertel, forest clearings, 500-600 m, 22.03.1995, K. 81.

*V. grandiflora* Scop. var. *grandiflora*  
Güvercinlik, 500-600 m, 08.04.1995, K. 76.  
Medit. & Euro.-Sib. element.

*V. sativa* L. subsp. *sativa*  
Bulutoğlu to Güvercinlik, fields, 500-600 m, 20.04.1995, K. 18.

*Lens orientalis* (Boiss.) Hand.-Mazz.  
Güvercinlik, riverside, 500-600 m, 20.04.1995, K. 21.

*Lathyrus cicera* L.  
Koçlar to Kamışlı, hills, 900-1000 m, 03.05.1995, K. 33.

*L. sativus* L.  
Kertel to Güvercinlik, landslide places, 500-600 m, 20.04.1995, K. 27.

*L. lycicus* Boiss.  
Kertel to Güvercinlik, 500-600 m, 27.04.1995, K. 51.  
Endemic, Medit. element.

*L. aphaca* L. var. *biflorus* Post  
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.04.1995, K. 119.

*Ononis natrix* L. subsp. *natrix*  
Kılavuzlu to Güvercinlik, 500-600 m, 03.11.1994, K. 117.  
Medit. element.

*O. spinosa* L. subsp. *leiosperma* (Boiss.) Sirj.  
Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 3.

*Trifolium boissieri* Guss. ex Boiss.  
Kılavuzlu, landslide places, 500-600 m, 25.02.1995, K. 114.  
Medit. element.

*T. erubescens* Fenzl  
N of Kertel, riverside, 500-600 m, 20.04.1995, K. 4.  
Medit. element.

*T. campestre* Schreb.  
Kılavuzlu, 500-600 m, 20.04.1995, K. 108.

*T. spumosum* L.  
Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 27.04.1995, K. 40.  
Medit. element.

*T. stellatum* L. var. *stellatum*  
Bulutoğlu, 500-600 m, riverside, 20.04.1995, K. 75.

*T. lucanium* Gasp.  
Güvercinlik, 500-600 m, 20.05.1995, K. 99.

*T. lappaceum* L.  
Kertel to Güvercinlik, 500-600 m, 7.04.1995, K. 39.  
Medit. element.

*T. purpureum* Lois. var. *purpureum*  
Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 20.05.1995, K. 103.

*T. scutatum* Boiss.  
Kılavuzlu, 500-600 m, 08.04.1994, K. 74.  
Medit. element.

*T. pauciflorum* d'Urv.  
Güvercinlik, 500-600 m, 20.05.1995, K. 101.  
Medit. element.

*Melilotus officinalis* (L.) Desr.  
Kılavuzlu to Bulutoğlu, 500-600 m, 20.05.1995, K. 94.

*M. alba* Desr.  
Ceyhan HES, riverside, 500-600 m, 03.11.1994, K. 116.

*Trigonella plicata* (Boiss. & P.W.Ball) Boiss.  
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 11.03.1995, K. 104.  
Endemic, Medit. element.

<i>T. mesopotamica</i> Hub.-Mor.	<i>Anthyllis tetraphylla</i> L.	28-ROSACEAE
Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 20.04.1995, K. 15/a.	S of Kertel, riverside, 500-600 m, 06.04.1995, K. 110.	<i>Cerasus microcarpa</i> (C.A.Mey.) Boiss. subsp. <i>tortuosa</i> (Boiss. & Hausskn.) Browicz
Ir.-Tur. element.	Medit. element.	Sarıçukur, roadside, 600-700 m, 17.06.1995, K. 528.
<i>T. kotschyi</i> Fenzl	<i>Coronilla emerus</i> L. subsp. <i>emeroides</i> (Boiss. & Spruner) Hrabétova	Ir.-Tur. element.
Kertel, landslide places, 500-600 m, 27.04.1995, K.42.	Kılavuzlu, landslide places, 500-600 m, 08.04.1995, K. 91.	<i>C. avium</i> (L.) Moench
Endemic, Ir.-Tur. element.	<i>Coronilla scorpioides</i> (L.) K.Koch	<i>Amygdalus orientalis</i> Mill.
<i>T. filipes</i> Boiss.	Güvercinlik, roadside, 500-600 m, 20.04.1995, K. 30.	Kılavuzlu, riverside, 500-600 m, 1994, K. 522.
Bulutoğlu, riverside, 500-600 m, 20.04.1995, K. 13.	<i>C. cretica</i> L.	<i>Potentilla reptans</i> L.
Ir.-Tur. element.	Kılavuzlu, riverside, 500-600 m, 08.04.1995, K. 73.	S of Bulutoğlu, riverside, 500-600 m, 20.04.1995, K. 520.
<i>T. crassipes</i> Boiss.	Medit. element.	<i>Sanguisorba minor</i> Scop. subsp. <i>muricata</i> (Spach) Briq.
Ceyhan HES, 500-600 m, 20.04.1995, K. 15/b.	<i>C. parviflora</i> Willd.	Kılavuzlu, riverside, 500-600 m, 20.04.1995, K. 530.
Ir.-Tur. element.	Kertel to Güvercinlik, riverside, 500-600 m, 27.04.1995, K. 53.	<i>Rosa foetida</i> Herrm.
<i>T. spicata</i> Sibth. & Sm.	Medit. element.	Kılavuzlu, 500-600 m, 27.04.1995, K. 529.
Kısık, riverside, 1300-1400 m, 03.05.1995, K. 15.	<i>C. grandiflora</i> Boiss.	Ir.-Tur. element.
Medit. element.	Kertel to Güvercinlik, riverside, 500-600 m, 27.04.1995, K. 44.	<i>R. canina</i> L.
<i>Medicago x varia</i> Martyn	Endemic, Medit. element.	Ceyhan HES, riverside, 500-600 m, 03.11.1994, K. 521.
Güvercinlik, riverside, 500-600 m, 12.06.1995, K. 2.	<i>Scorpiurus muricatus</i> L. var. <i>subvillosus</i> (L.) Fiori	<i>Crataegus orientalis</i> Pal. ex M.Bieb. var. <i>orientalis</i>
<i>M. minima</i> (L.) Bartal. var. <i>minima</i>	Ceyhan HES to Güvercinlik, 500-600 m, 20.05.1995, K. 100.	Kertel, riverside, 500-600 m, 22.03.1995, K. 526.
S of Kertel, riverside, 500-600 m, 08.04.1995, K. 58.	Medit. element.	<i>Pyrus syriaca</i> Boiss. var. <i>syriaca</i>
<i>M. rigidula</i> (L.) All. var. <i>cinerascens</i> (Jord.) Rouy	<i>Onobrychis aequidentata</i> (Sibth. & Sm.) d'Urv.	Kertel, streamside, 500-600 m, 22.03.1995, K. 526.
Kertel, riverside, 500-600 m, 27.04.1995, K. 47.	Hacınınoğlu, riverside, 500-600 m, 03.05.1995, K. 35.	29-LYTHRACEAE
<i>M. turbinata</i> (L.) All. var. <i>chiotica</i> Urb.	Medit. element.	<i>Lythrum salicaria</i> L.
Güvercinlik, riverside, 500-600 m, 27.04.1995, K. 49.	<i>O. armena</i> Boiss. & A.Huet	Ceyhan HES, riverside, 500-600 m, 06.07.1995, K. 703.
<i>Dorycnium pentaphyllum</i> Scop. subsp. <i>haussknechtii</i> (Boiss.) Gams	Kılavuzlu, fields, 500-600 m, 27.04.1995, K. 92.	30-CUCURBITACEAE
Menzelet to Kamişlı, riverside, 800-1000 m, 17.06.1995, K. 120.	<i>O. tournefortii</i> (Willd.) Desv.	<i>Cucumis melo</i> L.
Endemic, Ir.-Tur. element.	Sarıçukur, 600-700 m, 17.06.1995, K. 123.	Kertel, fields, 500-600 m, 05.06.1995, K. 761.
<i>Lotus angustissimus</i> L.	Endemic.	31-CRASSULACEAE
Ceyhan HES to Güvercinlik, riverside, 500-600 m, 20.07.1995, K. 687.	<i>Alhagi pseudalhagi</i> (M.Bieb.) Desv.	<i>Sedum caespitosum</i> (Cav.) DC.
<i>Hymenocarpus circinnatus</i> (L.) Savi	S of Kertel, riverside, 500-600 m, 17.06.1995, K. 121.	Kılavuzlu to Güvercinlik, rocky places, 500-600 m, 20.04.1995, K. 531.
Ceyhan HES, roadside, 500-600 m, 08.04.1995, K. 60.	Ir.-Tur. element.	Medit. element.

<i>Telmissa microcarpa</i> (Sm.) Boiss.	<i>Torilis arvensis</i> (Huds.) Link subsp. <i>arvensis</i>	<i>Inula graveolens</i> (L.) Desf.
Sarıçukur, riverside, 600-700 m, 27.04.1995, K. 593.	Ceyhan HES, riverside, 500-600 m, 12.06.1995, K. 284.	Sarıçukur, riverside, 800-900 m, 17.06.1995, K.202.
<b>32-APIACEAE</b>	<i>Caucalis platycarpus</i> L.	Medit. element.
<i>Eryngium campestre</i> L. var. <i>virens</i> Link	Koçlar to Kisik, fields, 1000-1100 m, 03.05.1995, K. 271.	<i>Helichrysum sanguineum</i> (L.) Kostel.
Ceyhan HES, riverside, 06.07.1995, K. 705.	<i>Daucus carota</i> L.	Kılavuzlu, riverside, 500-600 m, 20.05.1995, K. 172.
<i>Scandix stellata</i> Banks & Sol.	Güvercinlik, riverside, 500-600 m, 08.08.1995, K. 601.	Medit. element.
Güvercinlik, stony place, 500-600 m, 08.04.1995, K. 293.	<i>Artemia squamata</i> L.	<i>H. plicatum</i> DC. subsp. <i>plicatum</i>
<i>S. pecten-</i> <i>veneris</i> L.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, K. 288.	Kılavuzlu, landslide places, 500-600 m, 10.06.1995, K. 200.
S of Kertel, riverside, 500-600 m, 08.04.1995, K. 294.	<b>33-CAPRIFOLIACEAE</b>	<i>Filago eriocephala</i> Guss.
<i>Bifora testiculata</i> (L.) Spreng. ex Schult.	<i>Sambucus nigra</i> L.	Kamışlı, riverside, 900-1000 m, 03.05.1995, K. 158.
Bulutoğlu, riverside, 500-600 m, 12.06.1995, K. 281.	Ceyhan HES, riverside, 500-600 m, 12.06.1995, K. 747.	Medit. element.
<i>Bunium paucifolium</i> DC. var. <i>paucifolium</i>	Euro.-Sib. element.	<i>Logfia arvensis</i> (L.) Holub
Kertel, riverside, 500-600 m, 27.04.1995, K. 278.	<b>34-VALERIANACEAE</b>	Çakırdere, riverside, 800-900 m, 18.03.1995, K. 722.
Ir.-Tur. element.	<i>Valerianella vesicaria</i> (L.) Moench	<i>Aster alpinus</i> L.
<i>Pimpinella corymbosa</i> Boiss.	Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 580.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 05.06.1995, K. 127.
Ceyhan HES, fields, 500-600 m, 12.06.1995, K. 285.	<b>35-DIPSACACEAE</b>	<i>Erigeron acer</i> L. subsp. <i>acer</i>
Ir.-Tur. element.	<i>Cephalaria syriaca</i> (L.) Schrad.	Kılavuzlu to Güvercinlik, roadside, 500-600 m, 11.03.1994, K. 142.
<i>Heptaptera anisoptera</i> (DC.) Tutin	Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 764.	<i>Bellis perennis</i> L.
Ceyhan HES to Güvercinlik, fields, 500-600 m, 12.06.1995, K. 280.	<i>Scabiosa calocephala</i> Boiss.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 11.03.1995, K. 168.
<i>Bupleurum rotundifolium</i> L.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.05.1995, K. 719.	Euro.-Sib. element.
Sarıçukur, 800-1000 m, fields, 17.06.1995, K. 772.	Ir.-Tur. element.	<i>B. sylvestris</i> Cirillo
<i>B. gracile</i> d'Urv.	<i>Pterocephalus plumosus</i> (L.) Coult.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 11.03.1994, K. 141.
Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 269.	Kılavuzlu, landslide places, 500-600 m, 27.04.1995, K. 732.	Medit. element.
Medit. element.	<b>36-ASTERACEAE</b>	<i>Oronicum haussknechtii</i> Cavill.
<i>B. lycaonicum</i> Snogerup	<i>Helianthus annus</i> L.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.05.1995, K. 162.
Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 268.	Kılavuzlu to Güvercinlik, roadside, 500-600 m, 03.11.1995, K. 134.	Endemic.
Endemic, Medit. element.	<i>Xanthium spinosum</i> L.	<i>Senecio vulgaris</i> L.
<i>Ferulago cassia</i> Boiss.	Ceyhan HES, riverside, 500-600 m, 17.06.1995, K. 139.	S of Kertel, riverside, 500-600 m, 22.03.1995, K. 153.
Ceyhan Köprüsü, nr. Maraş, Hausskn.	<i>X. strumarium</i> L. subsp. <i>cavanillesii</i> (Schouw) D.Löve & Dans.	<i>Senecio vernalis</i> Waldst. & Kit Tan
Medit. element	Ceyhan HES, fields, 500-600 m, 18.06.1995, K. 140.	Güvercinlik, streamside, 500-600 m, 25.02.1995, K. 132.
<i>Tordylium syriacum</i> L.	<i>Calendula officinalis</i> L.	Güvercinlik, riverside, 500-600 m, 25.02.1995, K. 130.
Güvercinlik, riverside, 500-600 m, 08.04.1995, K. 291.		
Medit. element.		

<i>Anthemis hyalina</i> DC.	<i>C. virgata</i> Lam.	<i>Leontodon hispidus</i> L. var. <i>glabratus</i> (W.Koch.) Bisch.
Kılavuzlu to Güvercinlik, landslide places, 500-600 m, 22.03.1995, K. 149.	S of Kertel, landslide places, 500-600 m, 05.06.1995, K. 146.	Ceyhan HES to Güvercinlik, roadside, 500-600 m, 01.12.1994, K. 137.
<i>A. tricornis</i> Eig	<i>C. solstitialis</i> L. subsp. <i>soltstitialis</i>	Euro-Sib. element.
Ceyhan HES, riverside, 500-600 m, 08.04.1995, K. 188.	Kılavuzlu, 500-600 m, 12.06.1995, K. 196.	<i>Hedypnois cretica</i> (L.) Dum.-Cours.
Endemic, Medit. element.	<i>C. iberica</i> Trevir. ex Sprengel	Koçlar, riverside, 750-850 m, 03.05.1995, K. 180.
<i>A. haussknechtii</i> Boiss. & Reut.	Ceyhan HES, riverside, 500-600 m, 12.06.01995, K. 199.	Medit. element.
Çakırdere, riverside, 900-950 m, 02.05.1995, K. 164.	<i>C. urvillei</i> DC. subsp. <i>urvillei</i>	<i>Rhagadiolus stellatus</i> (L.) Gaertn. var. <i>edulis</i> (Gaertn.) DC.
<i>A. pungens</i> Yavin	Çukurçam, 1450-1500 m, 25.07.1995, K. 201.	Çakırdere, riverside, 700-800 m, K. 191.
Ceyhan river, 15 km W of Maraş, 500 m, D. 27459.	Medit. element.	<i>Sonchus asper</i> (L.) Hill. subsp. <i>glaucescens</i> (Jord.) P.W. Ball
Endemic.	<i>Crupina crupinastrum</i> (Moris) Vis.	Ceyhan HES, riverside, 500-600 m, 08.04.1995, K. 184.
<i>A. halophila</i> Boiss. & P.W.Ball	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.05.1995, K. 154.	<i>S. oleraceus</i> L.
Çakırdere to Koçlar, riverside, 900-1000 m, 03.05.1995, K. 157.	<i>Cnicus benedictus</i> L. var. <i>kotschy</i> Boiss.	S of Kertel district, landslide places, 500-600 m, 22.05.1995, K. 176.
Endemic, Medit. element.	Kılavuzlu, 500-600 m, 20.04.1995, K. 128.	<i>Taraxacum scaturiginosum</i> G. E. Haglund
<i>Achillea aleppica</i> DC. subsp. <i>aleppica</i>	<i>Xeranthemum longipapposum</i> M.A.Fisch. & C.A.Mey.	Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 08.04.1995, K. 186.
Sarıçukur, riverside, 600-700 m, 17.06.1995, K. 144.	Ceyhan HES, riverside, 500-600 m, 12.06.1995, K. 198.	<i>Chondrilla juncea</i> L. var. <i>juncea</i>
Ir.-Tur. element.	Ir.-Tur. element.	Ceyhan HES to Güvercinlik, riverside, 500-600 m, 20.07.1995, K. 680.
<i>Gundelia tournefortii</i> L. var. <i>tournefortii</i>	<i>Chardinia orientalis</i> (L.) Kuntze	<i>Crepis reuterana</i> Boiss. subsp. <i>reuterana</i>
Hacıoğlu, riverside, 1400-1500 m, 03.05.1995, K. 159.	Çakırdere, riverside, 900-1000 m, 03.05.1995, K. 155.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1995, K. 135.
<i>Cynara syriaca</i> Boiss.	Ir.-Tur. element.	Medit. element.
Güvercinlik, riverside, 500-600 m, 20.07.1995, K. 684.	<i>Scolymus hispanicus</i> L.	<i>C. sancta</i> (L.) Babc.
<i>Onopordum carduchorum</i> Bornm. & Beauverd	Koçlar, riverside, 1100-1200 m, 03.05.1995, K. 182.	Kılavuzlu, riverside, 500-600 m, 25.02.1995, K. 131.
Güvercinlik, riverside, 500-600 m, 05.06.1995, K. 170.	Medit. element.	<i>C. aspera</i> L.
Ir.-Tur. element.	<i>Cichorium endivia</i> L.	Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 08.04.1995, K. 187.
<i>Cirsium pubigerum</i> (Desf.) DC. var. <i>spinosum</i> Petr.	Kılavuzlu, riverside, 500-600 m, 03.11.1995, K. 136.	<b>37-CAMPANULACEAE</b>
Sarıçukur to Menzelet, riverside, 600-700 m, 17.06.1995, K. 145/b.	<i>Scorzonera mollis</i> M.Bieb. subsp. <i>szowitzzii</i> (DC.) D.F. Chamb.	<i>Michauxia campanuloides</i> L'Hér. ex Aiton
Ir.-Tur. element.	Baydemirli, riverside, 1400-1500 m, 18.03.1995, K. 173.	Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 646.
<i>C. elodes</i> M.Bieb.	Ir.-Tur. element.	Medit. element.
Bulutoğlu to Güvercinlik, riverside, 500-600 m, 20.04.1995, K. 129.	<i>S. kotschy</i> Boiss.	<i>Legousia speculum-veneris</i> (L.) Chaix
Ir.-Tur. element.	Ceyhan HES, roadside, 500-600 m, 22.03.1995, K. 17.	Ceyhan HES to Bulutoğlu, roadside, 500-600 m, 12.06.1995, K. 645.
<i>Centaurea aggregata</i> M.A.Fisch. & C.A.Mey. ex DC. subsp. <i>aggregata</i>	Ir.-Tur. element.	Medit. element.
Ceyhan HES, riverside, 500-600 m, 12.06.1995, K. 197.	<i>Tragopogon longirostris</i> Bisch. ex Sch. Bip. var. <i>longirostris</i>	
	Çakırdere, fields, 1100-1200 m, 03.05.1995, K. 173.	

<b>38-ERICACEAE</b>	<b>44-CONVOLVULACEAE</b>	<b>Onosma frutescens Lam.</b>
<i>Arbutus andrachne</i> L.	<i>Convolvulus arvensis</i> L.	Kılavuzlu, roadside, 500-600 m, 20.04.1995, K. 306.
Bulutoğlu, landslide places, under <i>Pinus brutia</i> forest, 500-600 m, 04.06.1995, K. 671. Medit. element.	Kılavuzlu, roadside, 500-600 m, 20.04.1995, K. 590.	Medit. element.
<b>39-PRIMULACEAE</b>	<i>Euro-Sib. element.</i>	<i>O. bourgaei</i> Boiss.
<i>Androsace maxima</i> L.	<i>C. germanicae</i> Boiss. & Hausskn.	Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 314.
Baydemirli, roadside, 1000-1200 m, 18.03.1995, K. 757.	Ceyhan HES, roadside, 500-600 m, 20.05.1995, K. 590.	Ir.-Tur. element.
<i>Anagallis arvensis</i> L. var. <i>caerulea</i> (L.) Gouan	Endemic, Ir.-Tur. element.	<i>Anchusa leptophylla</i> Roem. & Schult. subsp. <i>incana</i> (Ledeb.) D.F.Chamb.
Kısık, 1500-1600 m, 03.05.1995, K. 740.	<i>C. scammonia</i> L.	Kılavuzlu, riverside, 500-600 m, 11.03.1995, K. 297.
<b>40-EBENACEAE</b>	S of Kertel, landslide places, 500-600 m, 24.04.1995, K. 591.	Ir.-Tur. element.
<i>Diospyros kaki</i> L. f.	Medit. element.	<i>A. azurea</i> Miller var. <i>azurea</i>
Kertel, cultivated places, 500-600 m, 08.08.1995, K. 791.	<b>45-CUSCUTACEAE</b>	S of Kertel, landslide places, 500-600 m, 08.04.1995, K. 315.
<b>41-STYRACACEAE</b>	<i>Cuscuta planiflora</i> Ten.	<i>A. strigosa</i> Labill.
<i>Styrax officinalis</i> L.	Sarıçukur, field clearings, 700-800 m, 17.06.1995, K. 755.	Kılavuzlu, riverside, 500-600 m, 20.04.1995, K. 312.
Kılavuzlu to Bulutoğlu, roadside, 500-600 m, 08.08.1995, K. 641.	<b>46-BORAGINACEAE</b>	<i>Nonea melanocarpa</i> Boiss.
<b>42-OLEACEAE</b>	<i>Heliotropium haussknechtii</i> Bunge	Bulutoğlu, roadside, 500-600 m, 20.04.1995, K. 310.
<i>Jasminum fruticans</i> L.	Kılavuzlu to Kertel, fields, 500-600 m, 08.08.1995, K. 630.	Ir.-Tur. element.
Kılavuzlu, landslide places, 500-600 m, 05.06.1995, K. 647.	Endemic, Medit. element.	<i>N. ventricosa</i> (Sm.) Griseb.
Medit. element.	<i>Myosotis minutiflora</i> Boiss. & Reut.	Koçlar, riverside, 1200-1300 m, 03.05.1995, K. 304.
<i>Fontanesia philliraeoides</i> Labill. subsp. <i>philliraeoides</i>	Bulutoğlu, roadside, 500-600 m, 20.04.1995, K. 301.	Medit. element.
Kılavuzlu, landslide places, 500-600 m, 27.05.1995, K. 730.	Medit. element.	<b>47-SCROPHULARIACEAE</b>
Medit. element.	<i>Solenanthus stamineus</i> (Desf.) Wettst.	<i>Verbascum tauri</i> Boiss. & Kotschy
<i>Olea europaea</i> L. var. <i>europaea</i>	Ceyhan HES, riverside, 500-600 m, 20.04.1995, K. 309.	Kılavuzlu, 500-600 m, 20.04.1995, K. 587.
Kılavuzlu, fields, 500-600 m, 27.05.1995, K. 729.	<i>Buglossoides arvensis</i> (L.) I.M.Johnst.	Endemic, Medit. element.
Medit. element.	Kılavuzlu, riverside, 500-600 m, 08.04.1995, K. 727.	<i>V. tripolitanum</i> Boiss.
<b>43-GENTIANACEAE</b>	<i>Echium glomeratum</i> Poir.	S of Kılavuzlu, stony places, 500-600 m, K. 586.
<i>Centaurium erythraea</i> Rafn subsp. <i>erythraea</i>	E of Kertel, riverside, 500-600 m, 08.08.1995, K. 621.	Medit. element.
S of Kertel and Güvercinlik, riverside, 500- 600 m, 03.05.1995, K. 766.	<i>E. italicum</i> L.	<i>V. glomeratum</i> Boiss.
Euro-Sib. element.	Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 12.06.1995, K. 319.	Sarıçukur, roadside, 600-800 m, 17.06.1995, K. 583.
<i>C. spicatum</i> (L.) Fritsch	<i>E. angustifolium</i> Mill.	Ir.-Tur. element.
Bulutoğlu, roadside, 500-600 m, 12.06.1995, K. 748.	W of Cavgalar, riverside, 1000-1100 m, 17.06.1995, K. 750.	<i>V. lasianthum</i> Boiss. ex Benth.
	Medit. element.	Sarıçukur, roadside, 600-800 m, 17.06.1995, K. 582.

<i>V. anastasii</i> Nábelék	<i>O. heldreichii</i> (Reut.) Beck	Endemic, Medit. element.
Kılavuzlu, riverside, 500-600 m, 17.06.1995, K. 586.	Koçlar, fields, 900-1000 m, 27.04.1995, K. 784.	<i>Stachys annua</i> (L.) L. subsp. <i>annua</i> var. <i>lycaonica</i> R.Bhattacharjee
Endemic, Ir.-Tur. element.	Ir.-Tur. element.	Kılavuzlu, riverside, 500-600 m, 20.05.1995, K. 515.
<i>Scrophularia xanthoglossa</i> Boiss. var. <i>decipiens</i> (Boiss. & Kotschy) Boiss.	<b>49-ACANTHACEAE</b>	Ir.-Tur. element.
Ceyhan HES, roadside, 500-600 m, K. 540.	<i>Acanthus dioscoridis</i> L. var. <i>dioscoridis</i>	<i>Lallemantia peltata</i> (L.) M.A.Fisch. & C.A.Mey.
Ir.-Tur. element.	S of Kertel, riverside, 500-600 m, 05.06.1995, K. 507.	S of Kertel, landslide places, 500-600 m, 22.03.1995, K. 734.
<i>Anarrhinum orientale</i> Benth.	<b>50-VERBENACEAE</b>	Ir.-Tur. element.
Kılavuzlu, riverside, 500-600 m, 20.05.1995, K. 539.	<i>Verbena officinalis</i> L.	<i>Micromeria myrtifolia</i> Boiss. & Hohen.
Ir.-Tur. element.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 12.06.1995, K. 749.	Kertel, riverside, 500-600 m, 05.06.1995, K. 509.
<i>Linaria chalepensis</i> (L.) Mill. var. <i>chalepensis</i>	<i>Vitex agnus-castus</i> L.	Medit. element.
Bulutoğlu, roadside, 500-600 m, 20.05.1995, K. 563.	Kılavuzlu to Ceyhan HES, riverside, 500-600 m, 20.07.1995, K. 682.	<b>Thymbra spicata</b> L. var. <i>spicata</i>
Medit. element.	Medit. element.	Kertel, forest clearings, 500-600 m, 05.06.1995, K. 508.
<i>Linaria simplex</i> (Willd.) DC.	<b>51-LAMIACEAE</b>	Medit. element.
Ceyhan HES, riverside, 500-600 m, 20.05.1995, K. 544.	<i>Ajuga chamaepitys</i> (L.) Schreber. subsp. <i>laevigata</i> (Banks & Sol.) P.H.Davis	<i>Mentha longifolia</i> (L.) Huds. subsp. <i>typhoides</i> (Briq.) Harley var. <i>typhoides</i>
<i>Veronica bozakmanii</i> M.A.Fisch.	Çakırdere, riverside, 900-1000 m, 03.05.1995, K. 512.	Kılavuzlu, riverside, 500-600 m, 20.07.1995, K. 687.
Kertel, riverside, 500-600 m, 22.03.1995, K. 537.	Ir.-Tur. element.	<b>Lycopus europaeus</b> L.
<i>Veronica anagallis-aquatica</i> L.	<i>Teucrium multicaule</i> Montbret & Aucher ex Benth.	Bulutoğlu, riverside, 500-600 m, 03.11.1995, K. 502.
Kılavuzlu, riverside, 500-600 m, 20.05.1995, K. 752.	Çakırdere to Çobanlı, riverside, 950-1050 m, 03.05.1995, K. 759.	Euro-Sib. element.
<i>Veronica pectinata</i> L. var. <i>schizocalyx</i> (Freyen & Sint.) Bornm.	Ir.-Tur. element.	<i>Ziziphora capitata</i> L.
Sarıçukur, riverside, 600-800 m, 17.06.1995, K. 744.	<i>T. polium</i> L.	S of Kertel, stony places, 500-600 m, 27.04.1995, K. 513.
<i>Parentucellia latifolia</i> (L.) Caruel subsp. <i>latifolia</i>	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 12.06.1995, K. 745.	Ir.-Tur. element.
Kertel, landslide places, riverside, 500-600 m, 27.05.1995, K. 728.	<i>Phlomis pungens</i> Willd. var. <i>hispida</i> Hub.-Mor.	<i>Salvia suffruticosa</i> Montbret & Aucher ex Benth.
Medit. element.	Kertel, landslide places, 500-600 m, 05.06.1995, K. 88.	Valley of Ceyhan river, 15 km W of Maraş, D. 27440.
<b>48-OROBANCHACEAE</b>	<i>Lamium amplexicaule</i> L.	Ir.-Tur. element.
<i>Phelypaea coccinea</i> (M.Bieb.) Poir.	Ceyhan HES to Güvercinlik, riverside, 500-600 m, 25.02.1995, K. 501.	<i>S. multicaulis</i> Vahl
Kılavuzlu to Kertel, landslide places, 500-600 m, 27.04.1995, K. 672.	<i>Molucella laevis</i> L.	Çakırdere, 800-900 m, 03.05.1995, K. 511.
Ir.-Tur. element.	Sarıçukur, fields, 600-700 m, 17.06.1995, K. 771.	Ir.-Tur. element.
<i>Orobanche aegyptiaca</i> Pers.	Ir.-Tur. element.	<i>S. cryptantha</i> Montbret & Aucher ex Benth.
Kılavuzlu to Kertel, roadside, 500-600 m, 27.04.1995, K. 744.	<i>Stachys cretica</i> L. subsp. <i>mersinaea</i> (Boiss.) Rech. fil.	Kılavuzlu to Kertel, riverside, 500-600 m, 27.04.1995, K. 519.
	Sarıçukur, fields, 600-700 m, 17.06.1995, K. 506.	Endemic, Ir.-Tur. element.

<i>S. sclarea</i> L.	<i>E. falcata</i> L. subsp. <i>falcata</i> var. <i>falcata</i>	<i>Q. coccifera</i> L.
Bulutoğlu to Güvercinlik, riverside, 500-600 m, 20.04.1995, K. 503.	Kılavuzlu to Güvercinlik, roadside, 500-600 m, 20.07.1995, K. 675.	Kılavuzlu, landslide area, 500-600 m, 07.11.1995, K. 616.
<i>S. aethiopis</i> L.	<i>E. denticulata</i> Lam.	Medit. element.
Baydemirli, riverside, 1000-1100 m, 03.05.1995, K. 518.	Sarıçukur, roadside, 800-900 m, 17.06.1995, K. 457.	<b>60-SALICACEAE</b>
<i>S. cassia</i> Sam. ex Rech. fil.	Ir.-Tur. element.	<i>Salix pentandra</i> L.
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 20.05.1995, K. 514.	<i>E. forsskallii</i> (Boiss.) Gay	Güvercinlik, riverside, 500-600 m, 08.04.1995, K. 673.
Medit. element.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1995, K. 464.	Euro-Sib. element.
<b>52-PLANTAGINACEAE</b>	<b>54-URTICACEAE</b>	<i>Populus nigra</i> L. subsp. <i>nigra</i>
<i>Plantago major</i> L. subsp. <i>major</i>	<i>Parietaria judaica</i> L.	S of Kertel, cultivated places, 500-600 m, 27.04.1995, K. 790.
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 05.04.1995, K. 670.	Kılavuzlu to Ceyhan HES, rocky fields, 500-600 m, 12.06.1995, K. 741.	<b>61-RUBIACEAE</b>
<i>Plantago lanceolata</i> L.	<b>55-MORACEAE</b>	<i>Putoria calabrica</i> (L. f.) DC.
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 05.04.1995, K. 669.	<i>Morus nigra</i> L.	Kılavuzlu, roadside, 500-600 m, 11.03.1995, K. 566.
<b>53-EUPHORBIACEAE</b>	<i>S</i> of Kertel, cultivated places, 500-600 m, 10.06.1995, K. 536.	Medit. element.
<i>Chrozophora tinctoria</i> (L.) Raf.	<i>Ficus carica</i> L. subsp. <i>carica</i>	<i>Sherardia arvensis</i> L.
Kılavuzlu to Kertel, roadside, 500-600 m., 08.08.1995, K. 628.	Kılavuzlu to Güvercinlik, rocky fields, 500-600 m, 03.11.1995, K. 535.	Kılavuzlu to Ceyhan HES, roadside, 500-600 m, 11.03.1995, K. 576.
<i>Mercurialis annua</i> L.	<b>56-ULMACEAE</b>	Medit. element.
Güvercinlik, roadside, 500-600 m, 11.03.1995, K. 648.	<i>Ulmus minor</i> Mill. subsp. <i>canescens</i> (Melville) Browicz & Ziel.	<i>Crucianella macrostachya</i> Boiss.
<i>Euphorbia pubescens</i> Vahl	<i>S</i> of Kertel, 500-600 m, 08.08.1995, K. 788.	Kılavuzlu to Ceyhan HES, roadside, 500-600 m, 20.05.1995, K. 568.
Sarıçukur, fields, 600-800 m, 17.06.1995, K. 454.	Medit. element.	Medit. element.
<i>E. stricta</i> L.	<b>57-JUGLANDACEAE</b>	<i>C. imbricata</i> Boiss.
Kertel, cultivated places, 500-600 m, 20.07.1995, K. 458.	<i>Juglans regia</i> L.	Kertel, landslide places, 500-600 m, 05.06.1995, K. 736.
Euro-Sib. element.	<i>S</i> of Kertel, cultivated places, 500-600 m, 08.08.1995, K. 789.	Medit. element.
<i>E. helioscopia</i> L.	<b>58-PLATANACEAE</b>	<i>Asperula orientalis</i> Boiss. & Hohen.
Kılavuzlu to Kertel, riverside, 500-600 m, 27.04.1995, K. 461.	<i>Platanus orientalis</i> L.	<i>S</i> of Kertel, riverside, 500-600 m, 08.04.1995, K. 567.
<i>E. aleppica</i> L.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1995, K. 667.	Ir.-Tur. element.
Çakırdere, fields, 800-1000 m, 03.05.1995, K. 455.	<b>59-FAGACEAE</b>	<i>A. arvensis</i> L.
<i>E. szovitsii</i> M.A.Fisch. & C.A.Mey. var. <i>szovitsii</i>	<i>Quercus cerris</i> L. var. <i>cerris</i>	Çakırdere, riverside, 800-900 m, 03.05.1995, K. 571.
Kılavuzlu, roadside, 500-600 m, 10.06.1995, K. 465.	Kılavuzlu to Kertel villages, orchards, 500-600 m, 08.08.1995, K. 625.	Medit. element.
Ir.-Tur. element.	<i>Q. brantii</i> Lindl.	<i>Galium verum</i> L. subsp. <i>glabrescens</i> Ehrend.
	<i>S</i> of Kertel, cultivated places, 500-600 m, 08.08.1995, K. 616.	Kertel, fields, 500-600 m, 05.06.1995, K. 570.
	Ir.-Tur. element.	Ir.-Tur. element.

<i>G. cassium</i> Boiss.	<b>63-LILIACEAE</b>	<i>Gagea glacialis</i> K.Koch
Kertel, fields, 500-600 m, 05.06.1995, K. 263.	<i>Smilax aspera</i> L.	Baydemirli, meadows, 950-1050 m, 18.03.1995, K. 557.
Medit. element.	S of Kertel, riverside, 500-600 m, 08.08.1995, K. 623.	Ir.-Tur. element.
<i>G. spurium</i> L. subsp. <i>ibicinum</i> (Boiss. & Hausskn. ex Boiss.) Ehrend.	<i>Asphodelus aestivus</i> Brot.	<i>Colchicum stevenii</i> Kunth.
Baydemirli, field clearings, 900-1000 m, 18.03.1995, K. 575.	Kilavuzlu to Güvercinlik, riverside, 500-600 m, 11.03.1995, K. 548.	Kilavuzlu, landslide places, 500-600 m, 03.11.1995, K. 559.
Medit. element.	Medit. element.	Medit. element.
<i>G. tricornutum</i> Dandy	<i>Asphodeline damascena</i> (Boiss.) Baker subsp. <i>damascena</i>	<b>64-IRIDACEAE</b>
W of Kertel, field clearings, 500-600 m, 08.04.1995, K. 577.	Sarıçukur, riverside, 800-900 m, 17.06.1995, K. 770.	<i>Iris histrio</i> Rchb. f.
Medit. element.	Ir.-Tur. element.	Çakırdere, riverside, 900-1100 m, 18.03.1995, K. 560.
<i>G. haussknechtii</i> Ehrend.	<i>Allium flavum</i> L. subsp. <i>tauricum</i> (Besser ex Rchb.) Stearn var. <i>tauricum</i>	Medit. element.
Kertel, forest clearings, 500-600 m, 27.04.1995, K. 574.	Sarıçukur, roadside, 800-900 m, 17.06.1995, K. 551.	<i>Crocus graveolens</i> Boiss. & Reut.
Ir.-Tur. element.	Medit. element.	Kertel, under <i>Pinus brutia</i> forest, 500-600 m, 25.02.1995, K. 561.
<i>G. floribundum</i> Sm. subsp. <i>airoides</i> Hub.-Mor. ex Ehrend. & Schönb.-Tem.	<i>A. atroviolaceum</i> Boiss.	Medit. element.
Kilavuzlu to Güvercinlik, riverside, 500-600 m, 20.07.1995, K. 695.	Sarıçukur, fields, 800-900 m, 17.06.1995, K. 550.	<b>65-TYPHACEAE</b>
Endemic, Medit. element.	<i>A. cardiotermon</i> M.A.Fisch. & C.A.Mey.	<i>Typha latifolia</i> L.
<i>Callipeltis cucullaria</i> (L.) Steven	Sarıçukur, roadside, 800-900 m, 17.06.1995, K. 549.	Kilavuzlu to Güvercinlik, riverside, 500-600 m, 12.06.1995, K. 654.
Çakırdere to Koçlar, riverside, 900-1000 m, 03.05.1995, K. 573.	<i>Ornithogalum sphaerocarpum</i> A.Kern.	<i>Typha domingensis</i> Pers.
Ir.-Tur. element.	Kilavuzlu, fields, 500-600 m, 20.05.1995, K. 553.	Güvercinlik, riverside, 500-600 m, 06.07.1995, K. 696.
<i>Cruciata taurica</i> (Willd.) Ehrend.	<i>O. narbonense</i> L.	<b>66-JUNCACEAE</b>
Çakırdere, riverside, 900-1000 m, 03.05.1995, K. 572.	Bulutoğlu, riverside, 500-600 m, 20.04.1995, K. 554.	<i>Juncus inflexus</i> L.
Ir.-Tur. element.	Medit. element.	Kilavuzlu, riverside, 500-600 m, 03.11.1995, K. 428.
<i>Rubia tenuifolia</i> d'Urv. subsp. <i>brachypoda</i> (Boiss.) Ehrend. & Schönb.-Tem.	<i>Muscari neglectum</i> Guss.	<i>Juncus articulatus</i> L.
Kamışlı, riverside, 1000-1100 m, 03.05.1995, K. 569.	S of Kertel, under <i>Pinus brutia</i> forest, 18.03.1995, K. 556.	Kilavuzlu, riverside, 500-600 m, 03.11.1995, K. 424.
Medit. element.	<i>Bellevalia tauri</i> Feinbrun	Euro-Sib. element.
<b>MONOCOTYLEDONES</b>	Kilavuzlu, fields, 500-600 m, 20.04.1995, K. 555.	<b>67-CYPERACEAE</b>
<b>62-ARACEAE</b>	Endemic, Medit. element.	<i>Cyperus longus</i> L.
<i>Arum conophalloides</i> Koschy ex Schott var. <i>conophalloides</i>	<i>B. gracilis</i> Feinbrun	Güvercinlik, riverside, 500-600 m, 06.07.1995, K. 694.
Çakırdere, riverside, 900-1000 m, 03.05.1995, K. 534.	S of Kertel, under <i>Pinus brutia</i> forest, 500-600 m, 22.03.1995, K. 552.	<i>Schoenoplectus lacustris</i> (L.) Pal. subsp. <i>lacustris</i>
<i>Biarum carduchorum</i> (Schott) Engl.	Endemic, Ir.-Tur. element.	Bulutoğlu to Kilavuzlu, roadside, 500-600 m, 20.05.1995, K. 432.
Kilavuzlu to Bulutoğlu, roadside, 500-600 m, 07.11.1995, K. 787.		<i>Bolboschoenus maritimus</i> (L.) Pal. var. <i>maritimus</i>
Ir.-Tur. element.		S of Kertel, rocky places, 500-600 m, 08.08.1995, K. 635.

<i>Carex phyllostachys</i> C.A.Mey.	<i>B. sterilis</i> L.	<i>Vulpia ciliata</i> Dumort.
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 08.04.1995, K. 435.	Kılavuzlu, riverside, 500-600 m, 12.06.1995, K. 368.	Kertel, riverside, 500-600 m, 27.04.1995, K. 384.
Euro-Sib. element.	<i>B. riparius</i> Rehmann	<i>V. unilateralis</i> (L.) Stace
<i>C. otrubae</i> Podp.	Demirli, fields, 900-1000 m, 10.06.1995, K. 412.	Koçlar, field clearings, 700-800 m, 03.05.1995, K. 390.
Kılavuzlu to Güvercinlik, riverside, 500-600 m, 08.04.1995, K. 430.	<i>Avena sterilis</i> L. subsp. <i>Iudoviciana</i> (Durieu) Gillet & Magne	<i>Poa bulbosa</i> L.
Euro-Sib. element.	Güvercinlik, roadside, 500-600 m, 08.04.1995, K. 416.	Kılavuzlu, roadside, 500-600 m, 08.04.1995, K. 419.
<b>68-POACEAE</b>	Medit. element.	<i>Dactylis glomerata</i> L. subsp. <i>glomerata</i>
<i>Elymus lazicus</i> (Boiss.) Melderis. subsp. <i>divaricatus</i> (Boiss. & P.W.Ball) Melderis	<i>Trisetum thospiticum</i> Chrtek	Kılavuzlu, roadside, 500-600 m, 03.11.1994, K. 389.
Kısık, 800-950 m, 06.10.1995, K. 372.	Kılavuzlu, stony places, 500-600 m, 27.04.1995, K. 353.	Euro-Sib. element.
Endemic, Ir.-Tur. element.	Endemic, Ir.-Tur. element.	<i>Cynosurus echinatus</i> L.
<i>Aegilops umbellulata</i> Zhuk.	<i>T. turcicum</i> Chrtek	Kertel, riverside, 500-600 m, 08.08.1995, K. 607.
Kamışlı, 700-800 m, 03.05.1995, K. 400.	Kılavuzlu, sandy places, 500-600 m, 03.11.1995, K. 354.	Medit. element.
Ir.-Tur. element.	Ir.-Tur. element.	<i>Briza humilis</i> M.Bieb.
<i>A. triuncialis</i> L.	<i>Koeleria cristata</i> (L.) Pers.	S of Kertel, landslide places, 500-600 m, 27.04.1995, K. 383.
Çakırdere, 700-800 m, 10.06.1995, K. 75.	Kılavuzlu to Kertel, field clearings, 500-600 m, 27.04.1995, K. 389.	<i>Echinaria capitata</i> (L.) Desf.
<i>Hordeum murinum</i> L. subsp. <i>glaucum</i> (Steud.) Tzvelev	<i>Calamagrostis pseudophragmites</i> (Haller f.) Koeler	Kamışlı, riverside, 800-900 m, 03.05.1995, K. 401.
S of Kertel, riverside, 500-600 m, 08.04.1995, K. 418.	Güvercinlik, roadside, 500-600 m, 12.06.1995, K. 367.	<i>Melica persica</i> Kunth subsp. <i>inaequiglumis</i> (Boiss.) Bor
<i>Taeniatherum caput-medusae</i> (L.) Nevski subsp. <i>crinitum</i> (Schreber) Melderis	<i>Agrostis stolonifera</i> L.	Ceyhan Köprü, 15.7.1865, Hausskn.
Kertel, landslide places, 500-600 m, 08.04.1995, K. 371.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1994, K. 352.	<i>Phragmites australis</i> (Cav.) Trin. ex Steud.
Ir.-Tur. element.	<i>Phalaris arundinacea</i> L.	Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1995, K. 363.
<i>Bromus squarrosus</i> L.	Kılavuzlu, streamside, 500-600 m, 03.11.1994, K. 352.	<i>Setaria viridis</i> (L.) P.Beauv.
Çakırdere, riverside, 700-800 m, 10.06.1995, K. 376.	<i>Alopecurus myosuroides</i> Huds. var. <i>myosuroides</i>	S of Kertel, field clearings, 500-600 m, 27.04.1995, K. 349.
<i>Bromus scoparius</i> L.	Kılavuzlu, sandy places, 500-600 m, 20.05.1995, K. 406.	<i>Chrysopogon gryllus</i> (L.) Trin.
Koçlar, riverside, 750-800 m, 10.06.1995, K. 373.	Euro-Sib. element.	Güvercinlik, rocky places, 500-600 m, 08.08.1995, K. 393.
<i>B. tectorum</i> L.	<i>Phleum exaratum</i> Hochst. ex Griseb. subsp. <i>exaratum</i>	<i>Hyparrhenia hirta</i> (L.) Stapf
Sarıçukur, riverside, 600-700 m, 17.06.1995, K. 374.	Cavgalar, riverside, 800-900 m, 10.06.1995, K. 380.	Kılavuzlu, rocky places, 500-600 m, 20.05.1995, K. 350.

## Results and Discussion

As a result of the identification of the c. 700 plant specimens, 412 taxa belonging to 68 families and 262

genera were determined. Four taxa are from the *Pteridophyta* division, while the other 408 taxa belong to the *Spermatophyta*. Three taxa belong to *Gymnospermae*

and the other taxa are in the subdivision of *Angiospermae*. Three hundred and forty-eight taxa belong to the *Dicotyledones* and the remaining taxa belong to the *Monocotyledones*.

The number of endemic taxa in our area is 28 and the endemism rate is 6.7%. The endemism rate is 15.6% for Engizek Mountain (Duman, 1991) and 16.1% for Berit Mountain (Yıldız, 1984), i.e. higher than in our research area. The high altitude of the research area and the existence of local endemics in high places explain the higher endemism on Berit and Engizek mountains.

Among the endemic plants in our research area, *Convolvulus germanicae* and *Doronicum haussknechtii* are only known from Kahramanmaraş and they are local endemics for the B6 and C6 squares. The majority of the endemic plants collected from the research area, such as *Glaucium acutidentatum*, *Silene nuncupanda*, *Trigonella kotschyi*, *Anthemis tricornis*, *A. halophila*, *A. pungens*, *Heliotropium haussknechtii*, *Verbascum anastasii* and *Belleviala gracilis*, are localized around the Anatolian Diagonal (Ekim & Güner, 1986).

Of all the collected specimens, 54 taxa are new records for the C6 square [Davis (ed), 1965-1985, Davis et al. (eds), 1988; Donner, 1990; Duman & Aytaç, 1995; Duman & Karavelioğulları, 1995; Duman & Vural, 1991; Yıldız, 1984].

*Glycyrrhiza aspera* Pall., *Ferulago cassia* Boiss., *Anthemis pungens* Yavin, *Salvia suffruticosa* Montbret & Aucher ex Benth. and *Melica persica* Kunth, which we were unable to collect during this study, were collected by P.H. Davis, P.W. Ball and Hausskn, and so were added to the floristic list. Among these species, *Anthemis pungens* is an endemic and was gathered from the research area by P.H.Davis (P.H.Davis 1975).

*Euphorbia forsskalii* was determined to be a new record for the *Flora of Turkey* (Sümbül, 1991). It was known from C4 Antalya: Gazipaşa, c. 10 km along Gazipaşa-Anamur road, macchie shrub, 40-50 m, 27.7.1989, Sümbül 3409. This species was also collected during this study from Kılavuzlu to Güvercinlik, riverside, 500-600 m, 03.11.1995, K. 464. Therefore, the distribution of this species in Turkey is widened with our study.

The taxa, categorized according to phytogeographic region, may be given as follows: Mediterranean elements

with a rate of 23.5%, are most common, followed by Irano-Turanian elements, with a rate of 17.2%, and Euro-Siberian elements, with a rate of 3.4%. The remaining elements, with a rate of 55.9%, are multi-regional or of unknown phytogeographic origin.

The vast number of Mediterranean elements can be explained by the fact that the study area lies wholly within the Mediterranean phytogeographic region and at low altitude (500-1250 m). Moreover, our research area is near an urban area and it shows that the steppic plants belonging to Irano-Turanian elements do not spread at lower altitudes. Irano-Turanian elements are dominant, with rate of 23.4% on Berit Mountain and a rate of 35.5% on Engizek Mountain, which are near our study area. The dominance of the Irano-Turanian elements may be explained by the high altitude (700-3000 m) of these research areas.

Concerning the number of taxa, *Fabaceae* is the largest family in the study area, with 60 taxa. Following *Fabaceae*, *Asteraceae* is represented by 52 taxa, *Poaceae* by 31, *Brassicaceae* by 24 and *Lamiaceae* by 19. *Fabaceae* is also the most common family on Engizek Mountain. The family *Asteraceae* ranks first both in the *Flora of Turkey* and on Berit Mountain. In the research area, since *Fabaceae* is the largest family in terms of the number of taxa, it may be concluded that the area is parallel to the *Flora of Turkey* since *Fabaceae* is the second richest family in Turkey. *Fabaceae* and *Asteraceae* are the two richest families in the *Flora of Turkey*, in most floristic studies. This can be explained by the diversity of vegetation types, soil construction, climatic conditions and topography. Since the altitude is between 700 m and 3000 m for Berit and Engizek mountains, woodland, meadowland, steppe and rocky vegetation types are observed.

Concerning the number of taxa, the major genera in this study are as follows: *Trifolium* with 10, *Vicia* with 9, *Euphorbia* with 8, *Papaver* with 6, *Galium* with 6 and *Astragalus* with 4 taxa. The genera which have the highest number of taxa vary slightly between studies. The genus *Astragalus* ranks first on Berit and Engizek mountains, but in our study the genus *Trifolium* is first. One of the reasons for this is the ecological factors and habitat for *Astragalus*, which is known as a steppic plant of Irano-Turanian phytogeographic regions. Since our

research area is near settlements and the altitude is lower (500-1250 m), the ecological factors are not suitable for *Astragalus* and, therefore, it is in sixth position in this study. This irregularity can be explained by differences in the habitats of the study areas. For instance, although the genus *Astragalus* includes the highest number of species in Turkey and features prominently in steppe regions of Irano-Turanian phytogeographic regions, it is not well represented in agricultural areas or at lower altitudes.

## References

- Akman Y (1973). Aperçu Préliminaire Sur les Conditions Phytosociologiques de la Chaîne de l'Amanus Dans la Région du Hatay (I-II-III). *Com de la Fac-des Sci-d'Ank Serie C. Tome 17*.
- Akman Y (1990). *İklim ve Biyoiklim*. Ankara: Palme Yayınları. Mühendislik serisi:103.
- Çetik R (1981). Erciyes Dağı'nın (Kayseri) vejetasyonu. *SÜ Fen Fak Derg Sayı 2*: 23-37.
- Davis PH (ed.) (1965-1985). *Flora of Turkey and the East Aegean Islands*, Vol. 1-9. Edinburgh: Edinburgh Univ. Press.
- Davis PH, Mill, RR, Tan, K (eds) (1988). *Flora of Turkey and the East Aegean Islands(Supplement)*, Vol. 10. Edinburgh: Edinburgh Univ. Press.
- Donner J (1990). Distribution maps to P.H. Davis' "Flora of Turkey" 1-10. *Linzer Biol Beitr* 22(2): 381-515.
- Duman H (1991). Engizek Dağı Florası. *J Bot Fac Sci Arts Gazi Uni* 2: 57-96.
- Duman H (1995). Engizek Dağı (Kahramanmaraş) vejetasyonu. *Doğa TU Bot Derg* 19(2):179-213.
- Duman H, Aytaç Z (1995). New Floristic Records for the Grid Squares B6 and C6. *Tr J of Botany* 19(6): 615-621.
- Duman H, Karavelioğulları, F.A. (1995). New Floristic Records for the Grid Squares (A4, A8, B1, B3, B6, C1, C6). *Tr J of Botany* 19(6): 623-626.
- Duman H, Vural M (1991). C6 Karesinden Yeni Floristik kayıtlar. *Doğa TU Bot Derg* 15(2): 201-213.
- Düzenli A (1976). Hasan Dağı'nın Bitki Ekolojisi ve Bitki Sosyolojisi Yönünden Araştırılması. *Orn Arş Enst Derg* 22(2): 7-53.
- Ekim T, Güner A (1986). The Anatolian Diagonal: Fact or Fiction? *Proceedings of Royal Society of Edinburgh*, 89B: 69-77.
- Eyce B (1987). Melendiz Dağları (Niğde) Florasına Katkılar II. *SÜ Fen Edeb Fak Derg Sayı 6*: 111-118.
- Eyce B, Ocakverdi H (1987). Melendiz Dağları (Niğde) Florasına Katkılar I. *Doğa TU Bot Derg* 11(2): 241-255.
- Meteoroloji Bülteni (1994). T.C. Başbakanlık Devlet Meteoroloji İşleri Genel Müdürlüğü Araştırma ve Bilgi İşlem Dairesi Başkanlığı. Ankara.
- Quezel P (1973). Contribution à l'étude phytosociologique du massif de Tarsus. *Phytocoenology*, 1(2): 131-222.
- Quezel P, Pamukçuoğlu A (1973). Contribution à l'étude phytosociologique de quelques groupements forestiers du Taurus. *Fed Rep.* 84(3): 185-229.
- Sümbül, H (1991). Ten new species from Anatolia and two new records for the Flora of Turkey. *Edinb J Bot* 48(1): 27-40.
- Uslu T (1977). A Plant Ecological and Sociological Research on the Dune and on maquis vegetation between Mersin and Silifke. *Com de la Fac Sci D'Ank Serie C2 Tome 21 Supplement: 1*
- Vural M, Akman Y, Quezel P (1999). Contribution à l'étude de la végétation forestière du taurus central: analyse phyto-écologique d'un transect sud-nord, entre Silifke et Karaman. *Fitosociologia* 36(1): 3-21.
- Yıldız B (1984). C6 Karesi için yeni kayıtlar. *CÜ Fen-Edeb Fak Fen Bilimleri Derg* 2: 43-52.
- Zohary M (1973). Geobotanical Foundations of the Middle East Vol. 1-2. Stuttgart: Gustav Fischer Verlag.

Some vegetables and fruits, such as *Juglans regia*, *Cerasus avium*, *Vitis vinifera*, *Morus nigra* and *Ficus carica*, are cultivated in the research area.

## Acknowledgements

We are grateful to Prof. Dr. Mecit VURAL, Prof. Dr. Hayri DUMAN and Assoc. Prof. Dr. Zeki AYTAÇ for permission to use their facilities in GAZI Herbarium to identify plants during this study.